arXiv:1401.3636v1 [math.CV] 15 Jan 2014

On extensions of two results due to Ramanujan

Y. S. Kim[a], A. K. Rathie[b] and R. B. Paris[c]

[a] Department of Mathematics Education, Wonkwang University, Iksan, Korea
E-Mail: yspkim@wonkwang.ac.kr
[b] Department of Mathematics, Central University of Kerala,
Kasaragad 671328, Kerala, India
E-Mail: akrathie@gmail.com
[c] University of Abertay Dundee, Dundee DD1 1HG, UK
E-Mail: r.paris@abertay.ac.uk

Abstract

The aim in this note is to provide a generalization of an interesting entry in Ramanujan's Notebooks that relate sums involving the derivatives of a function $\phi(t)$ evaluated at 0 and 1. The generalization obtained is derived with the help of expressions for the sum of terminating ${}_3F_2$ hypergeometric functions of argument equal to 2, recently obtained in Kim *et al.* [*Two results for the terminating* ${}_3F_2(2)$ *with applications*, Bull. Korean Math. Soc. **49** (2012) pp. 621–633]. Several special cases are given. In addition we generalize a summation formula to include integral parameter differences.

Mathematics Subject Classification: 33C15, 33C20

Keywords: Hypergeometric series, Ramanujan's sum, Sums of Hermite polynomials

1. Introduction

Two of the many interesting results stated by Ramanujan in his Notebooks are the following theorems, which appear as Entries 8 [1, p. 51] and Entry 20 [1, p. 36], expressing an infinite sum of derivatives of a function $\phi(t)$ at the origin to another infinite sum of its derivatives evaluated at $t = 1$.

Entry 8. *Let $\phi(t)$ be analytic for $|t-1| < R$, where $R > 1$. Suppose that a and $\phi(t)$ are such that the order of summation in*

$$\sum_{k=0}^{\infty} \frac{2^k (a)_k}{(2a)_k \, k!} \sum_{n=k}^{\infty} \frac{(-1)^n}{n!} \, (-n)_k \phi^{(n)}(1)$$

may be inverted. Then

$$\sum_{k=0}^{\infty} \frac{2^k (a)_k \phi^{(k)}(0)}{(2a)_k \, k!} = \sum_{k=0}^{\infty} \frac{\phi^{(2k)}(1)}{2^{2k} (a+\frac{1}{2})_k \, k!}. \tag{1.1}$$

Entry 20. *Let $\phi(t) = \sum_{k=0}^{\infty} \phi^{(k)}(1)(t-1)^k/k!$ be analytic for $|t-1| < R$, where $R > 1$. Suppose that a and b are complex parameters such that the order of summation in*

$$\sum_{k=0}^{\infty} \frac{(a)_k}{(b)_k k!} \sum_{n=k}^{\infty} \frac{(-1)^n}{n!} (-n)_k \phi^{(n)}(1)$$

may be inverted. Then

$$\sum_{k=0}^{\infty} \frac{(a)_k\, \phi^{(k)}(0)}{(b)_k k!} = \sum_{k=0}^{\infty} \frac{(-1)^k (b-a)_k}{(b)_k k!} \phi^{(k)}(1). \tag{1.2}$$

Berndt [1] pointed out that Entry 8 can be established with the help of the results

$${}_2F_1\left[\begin{matrix}-2n, a \\ 2a\end{matrix}; 2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{1}{2})_n}, \qquad {}_2F_1\left[\begin{matrix}-2n-1, a \\ 2a\end{matrix}; 2\right] = 0$$

for non-negative integer n, where $(a)_n = \Gamma(a+n)/\Gamma(a)$ denotes the Pochhammer symbol.

In [4], each of the above theorems was generalized. The result in (1.1) was extended by replacing the denominatorial parameter $2a$ by $2a+j$, where $j = 0, \pm 1, \ldots, \pm 5$. The second result in (1.2) was extended by the inclusion of an additional pair of numeratorial and denominatorial parameters differing by unity to produce the following theorem.

Theorem 1. *Let $\phi(t)$ be analytic for $|t-1| < R$, where $R > 1$. Suppose that a, b, d and $\phi(t)$ are such that the order of summation in*

$$\sum_{k=0}^{\infty} \frac{(a)_k (d+1)_k}{(b)_k (d)_k k!} \sum_{n=k}^{\infty} \frac{(-1)^n}{n!} (-n)_k \phi^{(n)}(1)$$

may be inverted. Then

$$\sum_{k=0}^{\infty} \frac{(a)_k (d+1)_k}{(b)_k (d)_k} \frac{\phi^{(k)}(0)}{k!} = \sum_{k=0}^{\infty} \frac{(-1)^k (b-a-1)_k (f+1)_k}{(b)_k (f)_k} \frac{\phi^{(k)}(1)}{k!}, \tag{1.3}$$

where $f = d(b-a-1)/(d-a)$.

This result was extended to the case where a pair of numeratorial and denominatorial parameters differs by a positive integer m to produce

$$\sum_{k=0}^{\infty} \frac{(a)_k (d+m)_k}{(b)_k (d)_k} \frac{\phi^{(k)}(0)}{k!} = \sum_{k=0}^{\infty} \frac{(-1)^k (b-a-m)_k}{(b)_k} \frac{((\xi_m+1))_k}{((\xi_m))_k} \frac{\phi^{(k)}(1)}{k!},$$

where the $\xi_1, \ldots, \xi_m$ are the zeros of a certain polynomial of degree m.

In this note we shall similarly extend the result in (1.1) (when the parameter $2a$ is replaced by $2a+1$) by the inclusion of a pair of numeratorial and denominatorial parameters differing by unity. For this we shall require the summations of a ${}_3F_2$ hypergeometric function of argument equal to 2 obtained[1] in [3, Theorem 2]

$${}_3F_2\left[\begin{matrix}-2n, a, d+1 \\ 2a+1, d\end{matrix}; 2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{1}{2})_n}, \tag{1.4}$$

and

$${}_3F_2\left[\begin{matrix}-2n-1, a, d+1 \\ 2a+1, d\end{matrix}; 2\right] = \frac{(1-2a/d)}{2a+1} \frac{(\frac{3}{2})_n}{(a+\frac{3}{2})_n}, \tag{1.5}$$

for non-negative integer n. Several applications are presented in Section 3.

In the final section we generalize the result given in [1, p. 25] as Entry 9:

Entry 9. *If $Re\,(c-a) > 0$ then*

$$\sum_{k=1}^{\infty} \frac{(a)_k}{k(c)_k} = \psi(c) - \psi(c-a), \tag{1.6}$$

[1] It should be noted that the right-hand side of (1.4) is independent of the parameter d.

where $\psi(x)$ denotes the logarithmic derivative of $\Gamma(x)$.

We extend this summation to include additional numeratorial and denominatorial parameters differing by positive integers. To achieve this we make use of the generalized Karlsson-Minton summation formula for a ${}_{r+2}F_{r+1}$ hypergeometric function of unit argument.

2. Generalization of Ramanujan's result (1.1)

The result to be established in this section is given by the following theorem.

Theorem 2. *Let $\phi(t)$ be analytic for $|t-1|<R$, where $R>1$. Suppose that a, d and $\phi(t)$ are such that the order of summation in*

$$\sum_{k=0}^{\infty}\frac{2^k(a)_k(d+1)_k}{(2a+1)_k(d)_k\,k!}\sum_{n=k}^{\infty}\frac{(-1)^n}{n!}\,(-n)_k\phi^{(n)}(1)$$

may be inverted. Then

$$\sum_{k=0}^{\infty}\frac{2^k(a)_k(d+1)_k\phi^{(k)}(0)}{(2a+1)_k(d)_k\,k!}=\sum_{k=0}^{\infty}\frac{\phi^{(2k)}(1)}{2^{2k}(a+\frac{1}{2})_k\,k!}-\frac{(1-2a/d)}{2a+1}\sum_{k=0}^{\infty}\frac{\phi^{(2k+1)}(1)}{2^{2k}(a+\frac{3}{2})_k\,k!}. \tag{2.1}$$

Proof. Since $\phi(t)$ is analytic for $|t-1|<R$, we have

$$\phi^{(k)}(t)=\sum_{n=k}^{\infty}\frac{(-1)^n(-n)_k}{n!}\,\phi^{(n)}(1)$$

by suitable differentiation of the associated Taylor series. Then

$$\begin{aligned}S:=\sum_{k=0}^{\infty}\frac{2^k(a)_k(d+1)_k}{(2a+1)_k(d)_k}\,\frac{\phi^{(k)}(0)}{k!} &= \sum_{k=0}^{\infty}\frac{2^k(a)_k(d+1)_k}{(2a+1)_k(d)_k}\sum_{n=k}^{\infty}\frac{(-1)^n(-n)_k}{n!}\,\phi^{(n)}(1)\\ &= \sum_{n=0}^{\infty}\frac{(-1)^n\phi^{(n)}(1)}{n!}\sum_{k=0}^{n}\frac{2^k(a)_k(d+1)_k(-n)_k}{(2a+1)_k(d)_k k!}\\ &= \sum_{n=0}^{\infty}\frac{(-1)^n\phi^{(n)}(1)}{n!}\,{}_3F_2\left[\begin{matrix}-n,a,d+1\\2a+1,d\end{matrix};2\right]\end{aligned}$$

upon inversion of the order of summation by hypothesis.

If we now separate the above sum into terms involving even and odd n we obtain

$$S=\sum_{n=0}^{\infty}\frac{\phi^{(2n)}(1)}{(2n)!}\,{}_3F_2\left[\begin{matrix}-2n,a,d+1\\2a+1,d\end{matrix};2\right]-\sum_{n=0}^{\infty}\frac{\phi^{(2n+1)}(1)}{(2n+1)!}\,{}_3F_2\left[\begin{matrix}-2n-1,a,d+1\\2a+1,d\end{matrix};2\right].$$

Finally, using the summations in (1.4) and (1.5) and noting that $(2n)!=2^{2n}(\frac{1}{2})_n n!$, $(2n+1)!=2^{2n}(\frac{3}{2})_n n!$, we easily arrive at the right-hand side of (2.1). This completes the proof of the theorem. □

When $d=2a$ it is seen that (2.1) reduces to Ramanujan's result in (1.1).

3. Examples of Theorem 2

In this section, we provide some examples of different choices for the function $\phi(t)$ appearing in (2.1). Throughout this section we let k denote a non-negative integer.

(a) First we consider the simplest choice with $\phi(t) = \exp(xt)$, where x is an arbitrary variable (independent of t). Then $\phi^{(k)}(t) = x^k\,\phi(t)$ which satisfies the conditions for the validity of (2.1). Substitution of the derivatives into (2.1) and identification of the resulting series as hypergeometric functions immediately yields

$$e^{-x}{}_2F_2\left[\begin{matrix}a, d+1\\ 2a+1, d\end{matrix};2x\right] = {}_0F_1\left[\begin{matrix}-\\ a+\frac{1}{2}\end{matrix};\tfrac{1}{4}x^2\right] - \frac{(1-2a/d)x}{2a+1}\,{}_0F_1\left[\begin{matrix}-\\ a+\frac{3}{2}\end{matrix};\tfrac{1}{4}x^2\right]. \tag{3.1}$$

which is a result established by a different method in [9, Theorem 2]. In addition, it is interesting to observe that, since

$${}_0F_1\left[\begin{matrix}-\\ a+\frac{1}{2}\end{matrix};\tfrac{1}{4}x^2\right] = \Gamma(a+\tfrac{1}{2})(\tfrac{1}{2}x)^{\frac{1}{2}-a}\,I_{a-\frac{1}{2}}(x),$$

where I_ν is the modified Bessel function of the first kind, the result (3.1) can also be written in terms of I_ν.

(b) If we let $\phi(t) = \cosh(xt)$, we have $\phi^{(2k)}(t) = x^{2k}\cosh(xt)$ and $\phi^{(2k+1)}(t) = x^{2k+1}\sinh(xt)$. Then (2.1), after a little simplification making use of the identity

$$(a)_{2k} = (\tfrac{1}{2}a)_k(\tfrac{1}{2}a+\tfrac{1}{2})_k\,2^{2k},$$

and letting $d \to 2d$, reduces to

$${}_3F_4\left[\begin{matrix}\frac{1}{2}a, \frac{1}{2}a+\frac{1}{2}, d+1\\ \frac{1}{2}, a+\frac{1}{2}, a+1, d\end{matrix};x^2\right] = \cosh x\;{}_0F_1\left[\begin{matrix}-\\ a+\frac{1}{2}\end{matrix};\tfrac{1}{4}x^2\right] - \frac{(1-a/d)}{2a+1}\,x\sinh x\;{}_0F_1\left[\begin{matrix}-\\ a+\frac{3}{2}\end{matrix};\tfrac{1}{4}x^2\right]. \tag{3.2}$$

(c) If $\phi(t) = (x-t)^{-b}$, where b is an arbitrary parameter and $x > 2$, then

$$\phi^{(k)}(t) = \frac{(b)_k}{(x-t)^{b+k}}.$$

From (2.1), we therefore find

$$\left(\frac{x}{x-1}\right)^{-b}\sum_{k=0}^{\infty}\frac{2^k(a)_k(b)_k(d+1)_k}{(2a+1)_k(d)_k k!}\,x^{-k} = \sum_{k=0}^{\infty}\frac{2^{-2k}(b)_{2k}}{(a+\frac{1}{2})_k k!}\,(x-1)^{-2k}$$
$$-\frac{(1-2a/d)}{2a+1}\sum_{k=0}^{\infty}\frac{2^{-2k}(b)_{2k+1}}{(a+\frac{3}{2})_k k!}\,(x-1)^{-2k-1},$$

which yields

$$\left(\frac{x}{x-1}\right)^{-b}{}_3F_2\left[\begin{matrix}a, b, d+1\\ 2a+1, d\end{matrix};\frac{2}{x}\right] = {}_2F_1\left[\begin{matrix}\frac{1}{2}b, \frac{1}{2}b+\frac{1}{2}\\ a+\frac{1}{2}\end{matrix};\frac{1}{(x-1)^2}\right]$$
$$-\frac{(1-2a/d)b}{2a+1}(x-1)^{-1}{}_2F_1\left[\begin{matrix}\frac{1}{2}b+\frac{1}{2}, \frac{1}{2}b+1\\ a+\frac{3}{2}\end{matrix};\frac{1}{(x-1)^2}\right].$$

If we put $z = x/(1+x)$, this last result becomes

$$(1+z)^{-b}{}_3F_2\left[\begin{matrix}a, b, d+1\\ 2a+1, d\end{matrix};\frac{2z}{1+z}\right]$$
$$= {}_2F_1\left[\begin{matrix}\frac{1}{2}b, \frac{1}{2}b+\frac{1}{2}\\ a+\frac{1}{2}\end{matrix};z^2\right] - \frac{(1-2a/d)b}{2a+1}\,z\,{}_2F_1\left[\begin{matrix}\frac{1}{2}b+\frac{1}{2}, \frac{1}{2}b+1\\ a+\frac{3}{2}\end{matrix};z^2\right], \tag{3.3}$$

which has been obtained by different methods in [3, Theorem 3].

(d) Finally, with $\phi(t) = \exp(-x^2t^2/4)$, we have [8, p. 442]

$$\phi^{(k)}(t) = (-1)^k (x/2)^k e^{-x^2t^2/4}\, H_k(xt/2),$$

where H_k is the Hermite polynomial of order k. Since $H_{2k}(0) = (-1)^k(2k)!/k!$ and $H_{2k+1}(0) = 0$, it follows from (2.1) that

$$e^{x^2/4}\, {}_3F_3\left[\begin{matrix}\frac{1}{2}a, \frac{1}{2}a+\frac{1}{2}, d+1 \\ a+\frac{1}{2}, a+1, d\end{matrix}; -x^2\right]$$

$$= \sum_{k=0}^{\infty} \frac{(x/4)^{2k}}{(a+\frac{1}{2})_k k!}\, H_{2k}(x/2) + \frac{(1-a/d)}{a+\frac{1}{2}} \sum_{k=0}^{\infty} \frac{(x/4)^{2k+1}}{(a+\frac{3}{2})_k k!}\, H_{2k+1}(x/2) \qquad (3.4)$$

provided $a \neq -\frac{1}{2}, -\frac{3}{2}, \ldots$, where we have put $d \to 2d$.

The above series involving the Hermite polynomials can be expressed in terms of ${}_2F_2$ functions since [4, Eqs. (36), (37)]

$$\sum_{k=0}^{\infty} \frac{(x/4)^{2k}}{(a+\frac{1}{2})_k k!}\, H_{2k}(x/2) = e^{x^2/4}\, {}_2F_2\left[\begin{matrix}\frac{1}{2}a, \frac{1}{2}a+\frac{1}{2} \\ a, a+\frac{1}{2}\end{matrix}; -x^2\right],$$

$$\sum_{k=0}^{\infty} \frac{(x/4)^{2k}}{(a+\frac{3}{2})_k k!}\, H_{2k+1}(x/2) = x e^{x^2/4}\, {}_2F_2\left[\begin{matrix}\frac{1}{2}a+1, \frac{1}{2}a+\frac{3}{2} \\ a+\frac{3}{2}, a+2\end{matrix}; -x^2\right]$$

to yield

$${}_3F_3\left[\begin{matrix}\frac{1}{2}a, \frac{1}{2}a+\frac{1}{2}, d+1 \\ a+\frac{1}{2}, a+1, d\end{matrix}; -x^2\right] = {}_2F_2\left[\begin{matrix}\frac{1}{2}a, \frac{1}{2}a+\frac{1}{2} \\ a, a+\frac{1}{2}\end{matrix}; -x^2\right] + \frac{x^2(1-a/d)}{4a+2}\, {}_2F_2\left[\begin{matrix}\frac{1}{2}a+1, \frac{1}{2}a+\frac{3}{2} \\ a+\frac{3}{2}, a+2\end{matrix}; -x^2\right].$$

We remark that this last result can be derived alternatively by writing $(d+1)_k/(d)_k = 1 + k/d$ in the series expansion of the ${}_3F_3$ function combined with use of the result for contiguous ${}_2F_2$ functions [4]

$${}_2F_2\left[\begin{matrix}\alpha, \beta \\ \gamma, \delta\end{matrix}; z\right] - {}_2F_2\left[\begin{matrix}\alpha, \beta \\ \gamma, \delta+1\end{matrix}; z\right] = \frac{\alpha\beta z}{\gamma\delta(\delta+1)}\, {}_2F_2\left[\begin{matrix}\alpha+1, \beta+1 \\ \gamma+1, \delta+2\end{matrix}; z\right].$$

Finally, the representation (3.4) may be contrasted with the more general result obtained from Theorem 1 with $\phi(t) = \exp(-x^2t^2/4)$ given in [4, Eq. (40)] (with $x \to 2x$ and $d \to 2d$)

$$e^{x^2}\, {}_3F_3\left[\begin{matrix}\frac{1}{2}a, \frac{1}{2}a+\frac{1}{2}, d+1 \\ \frac{1}{2}b, \frac{1}{2}b+\frac{1}{2}, d\end{matrix}; -x^2\right] = \sum_{k=0}^{\infty} \frac{(b-a-1)_k (f+1)_k}{(b)_k (f)_k}\, x^k H_k(x),$$

where $f = 2d(b-a-1)/(2d-a)$.

4. Extension of the summation (1.6)

We employ the usual convention of writing the finite sequence of parameters $(a_1, \ldots, a_p)$ simply by (a_p) and the product of p Pochhammer symbols by $((a_p))_k \equiv (a_1)_k \ldots (a_p)_k$. In order to derive our extension of Ramanujan's sum (1.6) we make use of the generalized Karlsson-Minton summation theorem given below.

Theorem 3 *Let (m_r) be a sequence of positive integers and $m := m_1 + \cdots + m_r$. The generalized Karlsson-Minton summation theorem is given by [6, 7]*

$${}_{r+2}F_{r+1}\left[\begin{matrix}a, b, (d_r+m_r) \\ c, \quad (d_r)\end{matrix}; 1\right] = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} \sum_{k=0}^{m} \frac{(-1)^k (a)_k (b)_k C_{k,r}}{(1+a+b-c)_k} \qquad (4.1)$$

provided $\Re(c-a-b) > m$.

The coefficients $C_{k,r}$ appearing in (4.1) are defined for $0 \leq k \leq m$ by

$$C_{k,r} = \frac{1}{\Lambda} \sum_{j=k}^{m} \sigma_j \mathbf{S}_j^{(k)}, \qquad \Lambda = (d_1)_{m_1} \dots (d_r)_{m_r}, \tag{4.2}$$

with $C_{0,r} = 1$, $C_{m,r} = 1/\Lambda$. The $\mathbf{S}_j^{(k)}$ denote the Stirling numbers of the second kind and the σ_j $(0 \leq j \leq m)$ are generated by the relation

$$(d_1 + x)_{m_1} \cdots (d_r + x)_{m_r} = \sum_{j=0}^{m} \sigma_j x^j. \tag{4.3}$$

In [5], an alternative representation for the coefficients $C_{k,r}$ is given as the terminating hypergeometric series of unit argument

$$C_{k,r} = \frac{(-1)^k}{k!} \, {}_{r+1}F_r \left[\begin{matrix} -k, \ (d_r + m_r) \\ (d_r) \end{matrix} ; 1 \right]. \tag{4.4}$$

When $r = 1$, with $d_1 = d$, $m_1 = m$, Vandermonde's summation theorem [10, p. 243] can be used to show that

$$C_{k,1} = \binom{m}{k} \frac{1}{(d)_k}. \tag{4.5}$$

Our extension of Ramanujan's summation in (1.6) is given by the following theorem.

Theorem 4 *Let (m_r) be a sequence of positive integers and $m := m_1 + \cdots + m_r$. Then, provided $Re\,(c-a) > m$, we have*

$$\sum_{k=1}^{\infty} \frac{(a)_k \left((d_r + m_r)\right)_k}{(c)_k \left((d_r)\right)_k k} = \psi(c) - \psi(c-a) + \sum_{k=1}^{m} \frac{(-1)^k (a)_k \Gamma(k) C_{k,r}}{(1+a-c)_k}, \tag{4.6}$$

where $C_{k,r}$ are the coefficients defined in (4.3) and (4.4).

Proof. We follow the method of proof given in [1, p. 25]. If we differentiate logarithmically the left-hand side of (4.1) with respect to b and then set $b = 0$, making use of the simple fact that

$$\frac{d}{dx}(x)_k \bigg|_{x=0} = (k-1)!, \qquad k \geq 1,$$

we immediately obtain

$$\sum_{k=1}^{\infty} \frac{(a)_k \left((d_r + m_r)\right)_k}{(c)_k \left((d_r)\right)_k k}$$

when $\mathrm{Re}\,(c-a) > m$. Proceeding in a similar manner for the right-hand side of (4.1), we obtain

$$\psi(c) - \psi(c-a) + \sum_{k=1}^{m} \frac{(-1)^k (a)_k \Gamma(k) C_{k,r}}{(1+a-c)_k}.$$

This completes the proof of the theorem. □

When $r = 1$, the coefficients $C_{k,1}$ are given by (4.5) and we obtain the summation

$$\sum_{k=1}^{\infty} \frac{(a)_k (d+m)_k}{(c)_k (d)_k k} = \psi(c) - \psi(c-a) + m! \sum_{k=1}^{m} \frac{(-1)^k (a)_k}{k(m-k)!(1+a-c)_k (d)_k} \tag{4.7}$$

provided $\mathrm{Re}\,(c-a) > m$.

Acknowledgement: Y. S. Kim acknowledges the support of the Wonkwang University Research Fund (2014).

References

[1] B. C. Berndt, *Ramanujan's Notebooks*, Part II, Springer-Verlag, New York, 1987.

[2] J. Choi and A.K. Rathie, Two formulas contiguous to a quadratic transformation due to Kummer with an application, Hacettepe J. Math. Sci. **40(6)** (2011) 885–894.

[3] Y. S. Kim, J. Choi and A. K. Rathie, Two results for the terminating ${}_3F_2(2)$ with applications, Bull. Korean Math. Soc. **49(3)** (2012) 621–633.

[4] Y. S. Kim, A. K. Rathie and R. B. Paris, Generalization of two theorems due to Ramanujan, Integr. Trans. Spec. Funct. **24** (2012) 314–323.

[5] A. R. Miller and R. B. Paris, *On a result related to transformations and summations of generalized hypergeometric series*, Math. Communications **17**, 205–210, 2012.

[6] A. R. Miller and R. B. Paris, *Transformation formulas for the generalized hypergeometric function with integral parameter differences*, Rocky Mountain J. Math. **43**, 291–327, 2013.

[7] A. R. Miller and H. M. Srivastava, *Karlsson–Minton summation theorems for the generalized hypergeometric series of unit argument*, Integral Transforms and Special Functions **21**, 603–612, 2010.

[8] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clark (eds.), NIST Handbook of Mathematical Functions, Cambridge University Press, Cambridge, 2010.

[9] A. K. Rathie and T. K. Pogány, New summation formula for ${}_3F_2(\frac{1}{2})$ and a Kummer-type II transformation of ${}_2F_2(x)$, Math. Commun. **13** (2008) 63–66.

[10] L. J. Slater, Generalized Hypergeometric Functions, Cambridge University Press, Cambridge 1966.